

Study of CPO resonances on the intercombination line in ^{173}Yb

Pushpander Kumar, Alok K. Singh,* Vineet Bharti, and Vasant Natarajan
Department of Physics, Indian Institute of Science, Bangalore 560 012, India

Kanhaiya Pandey
Department of Physics, Indian Institute of Technology, Guwahati 781 039, India

Abstract

We study coherent population oscillations (CPO) in an odd isotope of the two-electron atom ^{173}Yb . The experiments are done using magnetic sublevels of the $F_g = 5/2 \rightarrow F_e = 3/2$ hyperfine transition of the $^1\text{S}_0 \rightarrow ^3\text{P}_1$ intercombination line. The experiments are done both with and without a magnetic field. In the absence of a magnetic field, the sublevels form a complex level structure that results in a saturated fluorescence spectrum with a linewidth that is larger than the 190 kHz natural linewidth of the transition. In the presence of a magnetic field (of magnitude 330 mG), a well-defined quantization axis is present which results in the formation of two M-type systems. The total fluorescence is then limited by spin coherence among the ground sublevels. In addition, the pump beam gets detuned from resonance which results in a reduced scattering rate from the $^3\text{P}_1$ state. Both of these result in a reduction of the linewidth to a subnatural value of about 100 kHz.

PACS numbers: 42.50.Gy, 42.50.Md, 32.70.Jz, 32.80.Qk

* Present address: Department of Physics of Complex Systems, Weizmann Institute of Science, Rehovot 7610001, Israel

I. INTRODUCTION

Coherent population oscillation (CPO) is a phenomenon in which two levels are coupled by two coherent electromagnetic fields of different amplitudes and frequencies [1]. This results in the population difference between the two levels to oscillate periodically at the beat frequency of the two fields. The linewidth of the resulting resonance is limited by the population relaxation rate, and can be subnatural for optical transitions. In recent times, it has received much interest from the quantum optics community [2–4] because it has important applications similar to the related phenomenon of coherent population trapping (CPT) [5]—like slowing and stopping of light [6, 7] and optical memories [8] for use in quantum information processing—but is more advantageous because of the following reasons.

- (i) There is no third level involved, so it can be observed in a larger class of materials including gases, liquids, and room temperature solids.
- (ii) It does not require a two-photon resonance, therefore laser jitter—both frequency and amplitude—becomes irrelevant.
- (iii) It offers spectral features that are broad, which implies that the medium will respond to very short optical pulses (of picosecond duration or less), whereas the shortest pulses that can be manipulated by CPT-based schemes are of nanosecond duration.

We have recently reported the first experimental observation of CPT and CPO in the two-electron atom Yb [9], a theoretical model for which was presented in Ref. [10]. The study was novel because it was an observation in a V-type system, a system which does not have such resonances in a one-electron atom. We used the intercombination line ($^1S_0 \rightarrow ^3P_1$ transition) at 556 nm and the even isotope ^{174}Yb . We used a small magnetic field (of order 0.5 G) to shift the CPT resonances away from line center, and retain only the CPO resonances. The linewidth of the central peak was then about 400 kHz, which was a factor of two higher than the natural linewidth of the upper $^3P_1$ state. This was because the peak was due to multiple CPO resonances, each of which had a resonant pump field.

Here, we report a study of CPO resonances in the odd isotope ^{173}Yb . The main difference is that the peak at line center does not move in the presence of a B field, which causes the pump beam to become detuned. The underlying reason is that the magnetic sublevels form many multilevel systems (with more than three levels) for which the two-photon CPT resonance condition does not change with detuning; unlike for a V-type system in the even isotope ^{174}Yb . The effect of the detuning is to cause a decrease in the decoherence due to spontaneous emission from the upper state, which causes the linewidth to reduce. This detuning also causes a decrease in the fluorescence signal, and an overall decrease in the signal-to-noise ratio (SNR). This explanation is borne out by a detailed density-matrix calculation of the levels involved.

We demonstrate this using the $F_g = 5/2 \rightarrow F_e = 3/2$ hyperfine transition, but the same trend of linewidth reduction is observed for the other two hyperfine transitions in this isotope. ^{173}Yb was preferred over the other odd isotope ^{171}Yb —which has only one transition—for the following reason. It gives us the ability to study transitions where the number of sublevels in the excited state is of three different kinds: smaller ($5/2 \rightarrow 3/2$), equal ($5/2 \rightarrow 5/2$), and larger ($5/2 \rightarrow 7/2$).

II. EXPERIMENTAL DETAILS

The experimental setup shown in Fig. 1 is the same as in our previous work in Ref. [9], and is reviewed here for completeness. Briefly, the $^1S_0 \rightarrow ^3P_1$ transition at 556 nm is generated by doubling the output of a fiber laser with linewidth of 70 kHz. The doubling is done in a commercial external cavity doubler using potassium niobate as the nonlinear crystal. As seen in the figure, part of the output beam from the doubler is used to lock the laser to the correct hyperfine transition. To achieve this, the laser beam is sent perpendicular to the atomic beam, and the resulting fluorescence is collected using a photomultiplier tube (PMT). The frequency dither required for generating the error signal is obtained by frequency modulating the acousto-optic modulator (AOM) in the path of the laser beam.




FIG. 1. Experimental setup. The probe beam is scanned by ± 4 MHz using a double-passed AOM. Figure key: $\lambda/2$ —halfwave retardation plate; PBS—polarizing beam splitter; PM fiber—polarizing maintaining fiber; PMT—photomultiplier tube.

The remaining part of the laser beam is used for the experiment. It is divided into two parts—one as a pump beam and the other as a probe beam—with orthogonal linear polarizations using a polarizing beam splitter (PBS). The pump beam is fixed in frequency, and is resonant with the transition used for locking the laser. This is achieved by using a

second AOM with an identical shift (of +100 MHz) as the one used for laser locking. The probe beam is scanned around this transition by double passing a third AOM at 50 ± 2 MHz. The double passing ensures directional stability as the AOM is scanned, and the variable shift of ± 2 MHz allows the probe beam to be scanned by ± 4 MHz around the resonance. The two beams are combined on another PBS, and then transported to the experimental chamber using a polarization maintaining (PM) fiber. The use of the fiber ensures perfect overlap of the two beams as the probe beam is scanned. The mode coming out of the fiber is perfectly Gaussian. The power in each beam is 40 μ W and its $1/e^2$ diameter is 5 mm, which gives a maximum Rabi frequency of 1.2Γ at beam center. The output of the fiber is sent perpendicular to the atomic beam, and the resulting fluorescence signal is detected using a second identical PMT.

The atomic beam used both for laser locking and the experiment is contained in an ultra high vacuum (UHV) chamber. The beam is generated by resistively heating a quartz ampule containing an ingot of unenriched Yb to a temperature of about 400°C. The vacuum chamber is maintained at a pressure below 10^{-8} torr using a 20 l/s ion pump.

The required magnetic field for the experiment is generated by winding three pairs of Helmholtz coils in the three orthogonal directions on the outside of the UHV chamber. The experiments with zero field are done by adjusting the currents in the three pairs of coils to get a symmetric spectrum, which tells us that all stray fields are nullified. A finite longitudinal field is generated by increasing the current in the pair of coils that are coaxial with the laser beam, and measured using a fluxgate magnetometer. The stability of the power supply driving the coils is better than 10^{-4} .

III. EXPERIMENTAL RESULTS

^{173}Yb has a nuclear spin I of $5/2$, hence there are three hyperfine transitions in this line: $5/2 \rightarrow 3/2$, $5/2 \rightarrow 5/2$, and $5/2 \rightarrow 7/2$. Experimental spectra for the first transition are shown in Fig. 2. The first thing to notice is that there is a single peak, both with and without the B field. The linewidth of the peak without a field—but with a resonant pump beam—is 270 kHz. This is slightly larger than the natural linewidth of the intercombination line of 190 kHz (determined by the lifetime of the $^3\text{P}_1$ state), and arises both because of multiple (CPT and CPO) resonances contributing to the peak and a saturated fluorescence effect [11]. In the presence of a B field of 330 mG, corresponding to a value of $\Delta_B = 139$ kHz, the linewidth reduces to a subnatural value of 100 kHz. This is because of reduced spontaneous emission from the upper state.

The same behavior, i.e. a linewidth reduction in the presence of a B field, is seen for the other two hyperfine transitions— $5/2 \rightarrow 5/2$ and $5/2 \rightarrow 7/2$. The results of linewidth measurements for all three cases are shown in Fig. 3. As seen, the linewidth of the resonance for the zero-field case shows a steady increase from the one for the $5/2 \rightarrow 3/2$ transition. This is because the number of sublevels increases steadily, which results in more and more resonances contributing to the peak.




FIG. 2. Experimental spectra for the $5/2 \rightarrow 3/2$ hyperfine transition, showing a single peak due to multiple CPO and CPT resonances. (a) With no field. (b) With a field of 330 mG.




FIG. 3. Linewidth of transitions with and without a B field. The error bars are the 1σ deviations obtained from the Lorentzian fit.

IV. THEORETICAL ANALYSIS

Since the probe and control beams are linearly polarized and the fluorescence in the perpendicular direction is collected, we have to consider transitions between magnetic sublevels coupled by σ^+ , σ^- , and π polarizations when no well-defined quantization axis is present, i.e. no B field is applied. However, when a B field is applied, transitions coupled by π polarization disappear because there is now a quantization axis.

The complete sublevel structure for the $5/2 \rightarrow 3/2$ transition with and without a B field is shown in Fig. 4. Transitions involving π polarization (no B field) are shown in part (a), while transitions without π polarization (with B field) are shown in part (b). The main thing to note from Fig. 4(a) is that no single three-level system is formed. For example, the $m_{F_g} = -3/2$ ground sublevel forms 2 three-level systems—a Λ -type system with the ground sublevel $m_{F_g} = -1/2$ and the excited sublevel $m_{F_e} = -3/2$, and a V-type system with two excited sublevels $m_{F_e} = -3/2$ and $m_{F_e} = -1/2$. The complicated multilevel structure implies that no dark resonance will be formed because any coherence between magnetic sublevels will be destroyed by $\Delta m = 0$ transitions. However, when a B field is applied, Fig. 4(b) shows that the transitions separate into two M-type systems. Each system can now be analyzed separately, and their contributions to the signal added.

The two sets of transitions are depicted in the figure with different colors. The first one (shown in red) involves the ground sublevels $m_{F_g} = -5/2, -1/2$, and $+3/2$. The second one (shown in blue) involves sublevels $m_{F_g} = -3/2, +1/2$, and $+5/2$. For ease of writing the Hamiltonian, we define the levels in the first system from $|1\rangle$ to $|5\rangle$. The Rabi frequency of the probe beam driving the $|i\rangle \rightarrow |j\rangle$ transition is Ω_{ij}^p , while its detuning is δ_{ij}^p . The corresponding parameters for the pump beam are Ω_{ij}^c and δ_{ij}^c . The Rabi frequencies of the probe and pump beams are weighted by the respective Clebsch-Gordan coefficients. In the rotating wave approximation, the Hamiltonian is

$$\begin{aligned}
H = & 0 |1\rangle \langle 1| \\
& - \hbar \delta_{12}^p |2\rangle \langle 2| \\
& - \hbar (\delta_{12}^p - \delta_{23}^p) |3\rangle \langle 3| \\
& - \hbar (\delta_{12}^p - \delta_{23}^p + \delta_{34}^p) |4\rangle \langle 4| \\
& - \hbar (\delta_{12}^p - \delta_{23}^p + \delta_{34}^p - \delta_{45}^p) |5\rangle \langle 5| \\
& + \left(\frac{\hbar \Omega_{12}^p}{2} + \frac{\hbar \Omega_{12}^c}{2} e^{-i\hbar(\delta_{12}^p - \delta_{12}^c)t} \right) |1\rangle \langle 2| \\
& + \left(\frac{\hbar \Omega_{23}^p}{2} + \frac{\hbar \Omega_{23}^c}{2} e^{i\hbar(\delta_{23}^p - \delta_{23}^c)t} \right) |2\rangle \langle 3| \\
& + \left(\frac{\hbar \Omega_{34}^p}{2} + \frac{\hbar \Omega_{34}^c}{2} e^{-i\hbar(\delta_{34}^p - \delta_{34}^c)t} \right) |3\rangle \langle 4| \\
& + \left(\frac{\hbar \Omega_{45}^p}{2} + \frac{\hbar \Omega_{45}^c}{2} e^{i\hbar(\delta_{45}^p - \delta_{45}^c)t} \right) |4\rangle \langle 5| \\
& + \text{Hermitean conjugate}
\end{aligned} \tag{1}$$

As seen from the figure, the probe detunings are such that

$$\delta_{12}^p - \delta_{23}^p = 0 \quad \text{and} \quad \delta_{12}^p - \delta_{23}^p + \delta_{34}^p - \delta_{45}^p = 0 \tag{2}$$




FIG. 4. Magnetic sublevels of the $5/2 \rightarrow 3/2$ hyperfine transition in ^{173}Yb . Transitions coupled by the pump beam are shown with solid lines, while those coupled by the probe beam are shown with dotted lines. The probe beam is scanned in frequency with a variable detuning δ . The pump beam has a fixed frequency resonant with the zero-field transition, and gets detuned in the presence of a B field.

since there is no splitting of the ground state. The equation of motion of the density matrix ρ is given by

$$\dot{\rho} = -\frac{i}{\hbar}[H, \rho] - \frac{1}{2}\{\Gamma, \rho\} \quad (3)$$

where Γ is the decay matrix. Its diagonal terms give the total decay rate (radiative and non-radiative) of the respective population terms. The off-diagonal terms represent the decoherence rate between states $|i\rangle$ and $|j\rangle$, such that

$$\Gamma_{ij} = \frac{\Gamma_{ii} + \Gamma_{jj}}{2} \quad (4)$$

Γ_{ij} between two ground states is the spin relaxation rate, and is taken to be $2\pi \times 10$ kHz.

The solution of Eq. (3) will be time dependent because the Hamiltonian is time dependent. Hence the solution can be written in the form

$$\rho_{ij} = \rho_{ij}^0 + \rho_{ij}^{\pm 1} e^{\pm i(\delta_{ij}^p - \delta_{ij}^c)t} + \rho_{ij}^{\pm 2} e^{\pm i(\delta_{ij}^p - \delta_{ij}^c)t} + \dots \quad (5)$$

ρ^0 —the DC component of ρ —is found by taking the Fourier transform of the numerical solution of ρ .

The fluorescence signal obtained in the experiment is proportional to the population in the upper levels, i.e. $\rho_{22} + \rho_{44}$. From the time dependence of this quantity shown in Fig. 5, we see that it oscillates around a mean value after a few lifetimes of the excited state. Thus, the system reaches “steady state” after this time.




FIG. 5. Transient behavior of $\rho_{22} + \rho_{44}$ (populations in upper levels) for the first M-type system shown in Fig. 4. Γ_e is the spontaneous decay rate of the excited state.

The other M-type system is analyzed in a similar manner. For clarity, it is denoted with prime symbols in Fig. 4(b). The “steady state” populations in the upper levels for the two M-type systems, i.e. $\rho_{22} + \rho_{44} + \rho_{2'2'} + \rho_{4'4'}$, is shown in Fig. 6. The main thing to notice is that there is only one peak, exactly as seen in the experimental spectrum. This is different from the spectrum for the even isotope, shown in Ref. [9], where four shifted peaks were seen in the presence of a B field. The linewidth of the peak is smaller than the natural linewidth because detuning will cause reduced decoherence from the upper state. But this comes at the price of reduced fluorescence signal, something which is also observed experimentally.




FIG. 6. Populations in the upper levels as a function of probe detuning for the two M-type systems formed using magnetic sub-levels of the $5/2 \rightarrow 3/2$ transition in the presence of B field of 330 mG. The Rabi frequencies of both the beams are taken to be at the maximum value from the experiment (1.2Γ).

V. CONCLUSIONS

In summary, we have studied CPO resonances in the odd isotope ^{173}Yb . The magnetic sublevel structure of the intercombination line used in this study is such that the peak at line center, which is due to both CPT and CPO resonances, does not split into multiple peaks in the presence of a B field; this is different from the even isotope used in our previous work [9]. The effect of a B field is to cause the pump beam to be detuned from resonance, which results in a reduced scattering rate from the excited state of the transition. The effect of this on the resonance peak is to reduce its linewidth. The linewidth reduces to a subnatural value of 100 kHz with a magnetic field of 330 mG. Such subnatural features could lead to new mechanisms for sub-Doppler cooling in divalent atoms compared to the more commonly used monovalent alkali-metal atoms [12].

ACKNOWLEDGMENTS

This work was supported by the Department of Science and Technology, India. P K and A K S acknowledge financial support from the Council of Scientific and Industrial Research, India; and V B from a D S Kothari post-doctoral fellowship of the University Grants Commission, India. The authors thank S Raghuveer for help with manuscript preparation.

-
- [1] M. S. Bigelow, N. N. Lepeshkin, and R. W. Boyd, Phys. Rev. Lett. **90**, 113903 (2003).
 - [2] T. Lauprêtre, S. Kumar, P. Berger, R. Faoro, R. Ghosh, F. Bretenaker, and F. Goldfarb, Phys. Rev. A **85**, 051805 (2012).
 - [3] K. T. Kapale, W. K. Lam, and G. S. Agarwal, in *The Rochester Conferences on Coherence and Quantum Optics and the Quantum Information and Measurement meeting* (Optical Society of America, 2013) p. M6.35.
 - [4] M.-A. Maynard, F. Bretenaker, and F. Goldfarb, Phys. Rev. A **90**, 061801 (2014).
 - [5] E. Arimondo, Prog. Optics **35**, 257 (1996).
 - [6] L. V. Hau, S. E. Harris, Z. Dutton, and C. H. Behroozi, Nature **397**, 594 (1999).
 - [7] O. Kocharovskaya, Y. Rostovtsev, and M. O. Scully, Phys. Rev. Lett. **86**, 628 (2001).
 - [8] A. Eilam, I. Azuri, A. V. Sharypov, A. D. Wilson-Gordon, and H. Friedmann, Opt. Lett. **35**, 772 (2010).
 - [9] A. K. Singh and V. Natarajan, New J. Phys. **17**, 033044 (2015).
 - [10] A. Vafafard, M. Mahmoudi, and G. S. Agarwal, Phys. Rev. A **93**, 033848 (2016).
 - [11] T. Yang, K. Pandey, M. S. Pramod, F. Leroux, C. C. Kwong, E. Hagiyeve, Z. Y. Chia, B. Fang, and D. Wilkowski, Eur. Phys. J. D **69**, 226 (2015).
 - [12] J. J. McFerran, J. Opt. Soc. Am. B **33**, 1278 (2016).